

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-97-1-0066

R&T Code 33e 1806

Dr. Richard S. Miller

Technical Report No. 100

COMPUTED PROPERTIES OF 1,1-DIAMINO-2,2-DINITROETHYLENE

by

Peter Politzer, M. Edward Grice, Monica C. Concha and Pat Lane

Department of Chemistry
University of New Orleans
New Orleans, LA 70148

August 15, 1997

DTIC QUALITY INSPECTED 2

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.






19970916 133

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE August 15, 1997	3. REPORT TYPE AND DATES COVERED Technical		
4. TITLE AND SUBTITLE Computed Properties of 1,1-Diamino-2,2-Dinitroethylene		5. FUNDING NUMBERS N00014-97-1-0066 Dr. Richard S. Miller R&T Code 33e 1806		
6. AUTHOR(S) Peter Politzer, M. Edward Grice, Monica C. Concha, and Pat Lane				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of New Orleans Department of Chemistry New Orleans, Louisiana 70148		8. PERFORMING ORGANIZATION REPORT NUMBER 100		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Code 333 800 N. Quincy Street Arlington, VA 22217		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Unlimited distribution.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) For 1,1-diamino-2,2-dinitroethylene, $(H_2N)_2C=C(NO_2)_2$, computational techniques have been used to predict its solid, liquid and gas phase heats of formation, its heats of vaporization and sublimation, its specific impulse, the C-NO ₂ and C-NH ₂ bond strengths, and its molecular geometry. There is significant intramolecular hydrogen bonding as well as electron delocalization, both of which stabilize the molecule and increase the C-NO ₂ and C-NH ₂ dissociation energies; this is expected to diminish the compound's impact and shock sensitivities.				
14. Subject Terms 1,1-diamino-2,2-dinitroethylene; $(H_2N)_2C=C(NO_2)_2$; heats of formation; heats of vaporization and sublimation; specific impulse; C-NO ₂ and C-NH ₂ bond dissociation energies; molecular geometry; intramolecular hydrogen bonding.			15. NUMBER OF PAGES 10	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Introduction:

1,1-Diamino-2,2-dinitroethylene, **1**, is of interest as a potentially significant energetic compound. It has the same molecular stoichiometry as HMX and RDX, and accordingly the same high value for moles of gaseous products (CO, N₂ and H₂O) per gram, 0.0405.

We have carried out a computational analysis of **1**, in order to provide data relevant to its potential energetic performance. These include: a) heats of formation (solid, liquid and gas phase), and heats of vaporization and sublimation; b) specific impulse; c) C–NO₂ and C–NH₂ bond strengths; and d) molecular geometry. In order to provide perspective in regard to the last point, we have also computed the geometries of the two isomers of **1** formed by interchanging NO₂ and NH₂ groups, **2** and **3**, as well as the two monosubstituted ethylenes, **4** and **5**.

**1****2****3****4****5**

Heats of formation:

We have used our density functional procedure [1] to compute the gas phase heat of formation of **1**. The vibrational energy was determined from the molecular stoichiometry [2]. The gas phase heat of formation is converted to the liquid and solid state values by subtracting, respectively, the heat of vaporization and the heat of sublimation. These are obtained by means of relationships that we have developed involving the computed electrostatic potential on the molecular surface [3,4].

Results:

$$\Delta H_f^{298K}(\text{gas}) = -1.2 \text{ kcal/mole} = -7.8 \text{ cal/g}$$

$$\Delta H_f^{298K}(\text{liquid}) = -16 \text{ kcal/mole} = -109 \text{ cal/g}$$

$$\Delta H_f^{298K}(\text{solid}) = -27 \text{ kcal/mole} = -183 \text{ cal/g}$$

$$\Delta H_{\text{vaporiz}} = 15 \text{ kcal/mole}$$

$$\Delta H_{\text{sublim}} = 26 \text{ kcal/mole}$$

It should be noted that the heats of formation for all three phases are negative. For comparison purposes, the solid phase heats of formation of HMX and RDX are, respectively, 60.4 cal/g and 76.1 cal/g [5].


Specific impulse:

The specific impulse of **1** was estimated using the Newpep program (Naval Weapons Center, China Lake, CA) and also a program that we have written [6]. Our calculated solid phase heat of formation was used as input data in both instances. The predicted specific impulse, relative to that of HMX, is:

Newpep:	0.93	Our program:	0.94
---------	------	--------------	------

C-NO₂ and C-NH₂ Bond Strengths:

The strengths of the C-NO₂ and C-NH₂ bonds in **1** were determined by computing their dissociation energies, i.e. $\Delta H(298 \text{ K})$ for the processes,



The geometries of **1** and the various products were optimized with the density functional B3P86/6-31+G** procedure [7], and the resulting energies were converted to enthalpies at 298 K using the calculated vibration frequencies [8].

The results are:

C-NO ₂ dissociation energy, (H ₂ N) ₂ C=C(NO ₂) ₂ :	$\Delta H(298 \text{ K}) = 70.0 \text{ kcal/mole}$
C-NH ₂ dissociation energy, (H ₂ N) ₂ C=C(NO ₂) ₂ :	$\Delta H(298 \text{ K}) = 111.6 \text{ kcal/mole}$

For comparison, the dissociation energies of the C-NO₂ and C-NH₂ bonds in H₃C-NO₂ and H₃C-NH₂, computed by the same procedure, are given below, together with the experimental values.


C-NO ₂ dissociation energy, H ₃ C-NO ₂ [9]:	59.8 kcal/mole
Experimental [10]:	60.8 kcal/mole
C-NH ₂ dissociation energy, H ₃ C-NH ₂ [9]:	84.6 kcal/mole
Experimental [11]:	84.9 kcal/mole

Thus the C-NO₂ and C-NH₂ bonds in **1** are stronger than in more typical cases such as H₃C-NO₂ and H₃C-NH₂.

Geometries:

In Figure 1 are presented the optimized geometries and corresponding energetic data, including relative enthalpies at 298 K, for **1** and for its isomers **2** and **3**. All calculations were at the density functional B3P86/6-31+G** level.

The most stable of these molecules is **1**, with **2** and **3** being higher in enthalpy at 298 K by 4 and 17 kcal/mole, respectively. Among the factors that determine these relative stabilities are (a) the possibilities for intramolecular hydrogen bonding, and (b) delocalization of electrons, as shown in the examples below.



(a) Intramolecular Hydrogen Bonding:

In Figure 1, the possible hydrogen bonds (which are primarily O---H in nature) are shown by dashed lines. There are two in **1**, and they can be considered to be relatively strong [12-14], with O---H distances of 1.77 Å. There are four in **2**, but they are not as strong, with O---H separations of about 1.90 Å and 2.08 Å. **3** also has four hydrogen bonds, one being N---H in nature, but they involve considerably greater distances and hence are relatively weak.

(b) *Electron Delocalization:*

The electron delocalization depicted above (sometimes termed “push-pull”) reflects the combined effects of donation by the amine group and withdrawal by the nitro group, with the latter enhancing the former. Structures **6B** and **7B** suggest that the extent to which this is occurring can be gauged by the shortening of the C–NH₂ and C–NO₂ bonds and the lengthening of the C=C. In order to permit such comparisons, Figure 1 includes the computed geometries of **4** and **5**, in which the combined effect is precluded.

Examination of the respective bond lengths shows that the degree of electron delocalization is greatest in **1** and least in **3**. In the latter, the effect of one of the NH₂, NO₂ pairs is quite similar to those in **2**; however the other NH₂, NO₂ pair is not interacting in this fashion, as can be seen from the rotated and pyramidal nature of the NH₂ (which permits additional hydrogen bonding).


Comments:

The analysis of molecular geometries has shown that **1** is significantly stabilized by both intramolecular hydrogen bonding and electron delocalization, to a greater extent than either **2** or **3**. Since both of these factors are diminished by removal of either an NO₂ or an NH₂ group, it follows that they should have the effect of increasing the C–NO₂ and C–NH₂ dissociation energies, as we have indeed found to be the case. In view of the widely-held belief that rupture of the C–NO₂ bond is a critical step in the decomposition processes of many energetic molecules [15-24], it can be anticipated that the greater strengths of these bonds in **1** will decrease its impact and shock sensitivities. On the other hand, the anomalous stability of **1** is responsible for its negative heats of formation.

References

1. D. Habibollahzadeh, M. E. Grice, M. C. Concha, J. S. Murray and P. Politzer, *J. Comp. Chem.* **16**, 654 (1995).
2. M. E. Grice and P. Politzer, *Chem. Phys. Lett.* **244**, 295 (1995).
3. J. S. Murray and P. Politzer, in *Quantitative Treatments of Solute/Solvent Interactions*, J. S. Murray and P. Politzer, Eds., Elsevier, Amsterdam, 1994, ch. 8.
4. P. Politzer, J. S. Murray, M. E. Grice, M. DeSalvo and E. Miller, *Mol. Phys.* **91**, 923 (1997).
5. J. Köhler and R. Meyer, *Explosives*, 4th ed., VCH Publishers, New York, 1993 .
6. P. Politzer, J. S. Murray, M. E. Grice and P. Sjöberg, in *Chemistry of Energetic Materials*, G. A. Olah and D. R. Squire, Eds., Academic Press, New York, 1991, ch. 4.
7. M. J. Frisch, G. W. Trucks, H. B. Schlegel, P. M. W. Gill, B. G. Johnson, M. A. Robb, J. R. Cheeseman, T. A. Keith, G. A. Petersson, J. A. Montgomery, K. Raghavachari, M. A. Al-Laham, V. G. Zakrzewski, J. V. Ortiz, J. B. Foresman, J. Cioslowski, B. B. Stefanov, A. Nanayakkara, M. Challacombe, C. Y. Peng, P. Y. Ayala, W. Chen, M. W. Wong, J. L. Andres, E. S. Replogle, R. Gomperts, R. L. Martin, D. J. Fox, J. S. Binkley, D. J. Defrees, J. Baker, J. P. Stewart, M. Head-Gordon, C. Gonzalez and J. A. Pople, Gaussian 94, (Gaussian, Inc., Pittsburgh, PA, 1995).
8. W. J. Hehre, L. Radom, P. v. R. Schleyer and J. A. Pople, *Ab Initio Molecular Orbital Theory*, Wiley-Interscience, New York, 1986 .
9. J. J. M. Wiener and P. Politzer, *J. Mol. Struct. (Theochem)* , in press.
10. D. R. Lide, Ed. *Handbook of Chemistry and Physics*, 71st ed., CRC Press, Boca Raton, FL, 1990.
11. D. F. McMillen and D. M. Golden, *Ann. Rev. Phys. Chem.* **33**, 493 (1982).
12. C. Ceccarelli, G. A. Jeffrey and R. Taylor, *J. Mol. Struct.* **70**, 255 (1981).
13. R. Taylor, *J. Mol. Struct.* **73**, 125 (1981).
14. A. Gavezzotti and G. Filippini, *J. Phys. Chem.* **98**, 4831 (1994).
15. A. Delpuech and J. Cherville, *Propell. Explos.* **3**, 169 (1978).
16. J. Sharma and F. J. Owens, *Chem. Phys. Lett.* **61**, 280 (1979).
17. M. J. Kamlet and H. G. Adolph, *Proceedings, Seventh Symposium (International) on Detonation*, Naval Surface Warfare Center, Report NSWCMP-82-334, Silver Springs, MD, 1981, p. 84.
18. J. Sharma, W. L. Garrett, F. J. Owens and V. L. Vogel, *J. Phys. Chem.* **86**, 1657 (1982).
19. A. C. Gonzalez, C. W. Larson, D. F. McMillen and D. M. Golden, *J. Phys. Chem.* **89**, 4809 (1985).
20. F. J. Owens, *J. Mol. Struct. (Theochem)* **121**, 213 (1985).

21. F. J. Owens and J. Sharma, *J. Appl. Phys.* **51**, 1494 (1985).
22. J. S. Murray and P. Politzer, in *Chemistry and Physics of Energetic Materials*, S. N. Bulusu, Ed., Kluwer, Dordrecht, The Netherlands, 1990, ch. 8.
23. P. Politzer and J. S. Murray, *Mol. Phys.* **86**, 251 (1995).
24. P. Politzer and J. S. Murray, *J. Mol. Struct.* **376**, 419 (1996).

Figure 1. B3P86/6-31+G** optimized geometries^a and energies.^b

$E = -599.70979$ hartrees

ZPE = 58.23 kcal/mole


$\Delta H(0 \rightarrow 298) = 64.58$ kcal/mole

Relative H(298 K) = 0

Sums of carbon bond angles = 360°

Sums of nitro bond angles = 360°

Sums of amine bond angles = 358°



$E = -599.70209$ hartrees

ZPE = 57.59 kcal/mole


$\Delta H(0 \rightarrow 298) = 64.08$ kcal/mole

Relative H(298 K) = 4.3 kcal/mole

Sums of carbon bond angles = 360°

Sums of nitro bond angles = 360°

Sums of amine bond angles = $357^\circ, 358^\circ$



$E = -599.68278$ hartrees

ZPE = 58.07 kcal/mole


$\Delta H(0 \rightarrow 298) = 64.80$ kcal/mole

Relative H(298 K) = 17.2 kcal/mole

Sums of carbon bond angles = 360°


Sums of nitro bond angles = 360°

Sums of amine bond angles = $331^\circ, 350^\circ$



Sums of carbon bond angles = 360°

Sums of amine bond angles = 346°



Sums of carbon bond angles = 360°

Sums of nitro bond angles = 360°

^aDistances are in Angstroms.

^b E = energy minimum at 0 K; it does not include the zero-point energy (ZPE). $\Delta H(0 \rightarrow 298)$ is the enthalpy change in going from 0 K to 298 K; it includes the ZPE. Since $E(0 \text{ K}) = H(0 \text{ K})$, it follows that $H(298 \text{ K}) = E(0 \text{ K}) + \Delta H(0 \rightarrow 298)$.